

The Distribution Route from Ancestors to Descendants

Baruch Fischer and Moshe Zakai
 Department of Electrical Engineering
 Technion, Haifa 32000, Israel

E-mail: fischer@ee.technion.ac.il , zakai@ee.technion.ac.il

Abstract

We study the distribution of descendants of a known personality, or of anybody else, as it propagates along generations from father or mother through any of their children. We ask for the ratio of the descendants to the total population and construct a model for the route of Distribution from Ancestors to Descendants (DAD). The population ratio r_n is found to be given by the recursive equation $r_{n+1} \approx (2 - r_n)r_n$, that provides the transition from the n -th to the $(n + 1)$ th generation. $r_0 = 1/N_0$ and N_0 is the total relevant population at the first generation. The number of generations it takes to make half the population descendants is $\log N_0 / \log 2$ and additional ~ 4 generations make everyone a descendent (=the full descendant spreading time). These results are independent of the population growth factor even if it changes along generations. As a running example we consider the offspring of King David. Assuming a population between $N_0 = 10^6$ and $5 \cdot 10^6$ of Israelites at King David's time (~ 1000 BC), it took 24 to 26 generations (about 600-650 years, when taking 25 years for a generation) to make every Israelite a King David descendent. The conclusion is that practically every Israelite living today (and in fact already at 350-400 BC), and probably also many others beyond them, are descendants of King David. We note that this work doesn't deal with any genetical aspect. We also didn't take into account here any geo-social-demographic factor. Nevertheless, along tens of generations, about 120 from King David's time till today, the DAD route is likely to govern the distribution in communities that are not very isolated.

1. Introduction

The well known Galton-Watson (GW) process [1, 2] investigates the extinction of surnames which propagate from father to son. We consider here a model which unlike the GW model depends on both parents, namely the offspring of a known personality which propagates through father or mother to their children and we ask for the ratio of the descendants to the total population. As an example we will consider the offspring of King David.

King David lived about 3,000 years ago. We assume that a generation (from birth until marriage and children) is 25 years, and each married couple has $2g$ children, where g is the growth factor per generation. Let N_n denote the number of Israelites at the n -th generation and $N_n = N_0 \cdot g^n$ where $n = 0, 1, 2, \dots, 120 (= 3000/25)$. Let $\tilde{D}_n$ denote the number of descendants (male and female) of King David at the n -th generation and $\tilde{C}_n = N_n - \tilde{D}_n$ denote the non-descendants. We start at the first generation with $\tilde{D}_0 = 1$, the dynasty founder. Our problem is to estimate the ratio $r_n = \tilde{D}_n/N_n$ after n generations. This will show, in particular, that practically all Israelites today are descendants. It is possible that a family disappears after a few generations (discussed in Section 6), but we assume throughout the paper that it doesn't happen. In our example with what we know about King David and his son Solomon, we do not have to worry about that. However if we are not sure about that we can say, as we discuss later, that *either all Israelites are his descendant or none*. Therefore if there is one descendant then all are descendants.

It will be shown that for a population of N_0 it took $\log N_0 / \log 2$ generations to make half the population descendants of King David. Additional four generations made all of them his descendants. That is the DAD full spreading time. The transitions region between low to high spreading ratio is very quick, a few generations. It is not only King David; the same relation exists regarding anyone else of his era (or any other early era) whose family survived in the first few generations (discussed in Section 6), including for example less admirable characters in the Bible like Nabal... Assuming a population between $N_0 = 10^6$ and $N_0 = 5 \cdot 10^6$ Israelites at King David's time (~ 1000 BC) [3], it took 24 to 26 generations (600-650 years, when taking 25 years for a generation) to ensure that every Israelite was his

descendant. That means that every Israelite living at 400 BC, the beginning of the era of the Second Temple in Jerusalem, was already a descendant of King David.

An interesting feature of the DAD route is that the descendant population ratio and the spreading time depend on N_0 but not on the population growth factor g even if g is generation dependent.

2. The rule of passing from $\tilde{D}_n$ to $\tilde{D}_{n+1}$

Two presentations of the same result are given.

The First Presentation: Let N_n be the total number of males and females at the n -th generation of a certain community, and among them $\tilde{D}_n$ (Davidian) descendants of the dynasty we follow. The number of males and females are assumed equal. Further assume that we have $N_n/2$ cards with all the female names, one name per card, and similarly in another box $N_n/2$ cards for all the males. Consider now a "match maker" picking up randomly and independently one card from each box, and combining them to a single card (marriage) with the two names on it. At this stage add to each of the two names information whether he (she) is or isn't a D-descendant. Throw away all cards where both names are non-descendants. The probability that a descendant married a descendant is $\tilde{D}_n/N_n$ and for a descendant marrying a non-descendant it is $(N_n - \tilde{D}_n)/N_n$.

The conditional expectation $E(\tilde{D}_{n+1}|\tilde{D}_n)$ (the number of descendants at the $(n+1)$ th generation conditioned on $\tilde{D}_n$, the number of descendants at the n -th generation) follows from the fact that for $g = 1$ a married couple who are both descendants will have 2 descendants. In this case the number of descendants will not change. On the other hand if a descendant married a non-descendant the one descendant in the n -th generation will generate two descendants for the $(n+1)$ th generation.

Therefore,

$$\begin{aligned}
E(\tilde{D}_{n+1}|\tilde{D}_n) &= g \cdot \left[\begin{array}{l} \text{the probability that} \\ \text{a descendant married} \\ \text{a descendant} \end{array} \right] \cdot \tilde{D}_n + 2g \cdot \left[\begin{array}{l} \text{the probability that} \\ \text{a descendant married} \\ \text{a non-descendant} \end{array} \right] \cdot \tilde{D}_n \\
&= \left(1 \cdot \frac{\tilde{D}_n}{N_n} + 2 \cdot \frac{N_n - \tilde{D}_n}{N_n} \right) \cdot g\tilde{D}_n = \left(2 - \frac{\tilde{D}_n}{N_n} \right) \cdot g\tilde{D}_n.
\end{aligned} \tag{2..1}$$

Note that $\tilde{D}_n$ and N_n include males and females, and we assume throughout the paper that their numbers are equal. We also comment that this statistical model can be viewed as a random walk process, as discussed below in section 3.

We can simplify the analysis by a simple renormalization procedure that shows right away that the DAD route is independent of the growth factor g . We normalize by setting:

$$D_n = \frac{\tilde{D}_n}{g^n}, \quad C_n = \frac{\tilde{C}_n}{g^n} = N_0 - D_n, \tag{2..2}$$

and then with $N_n = N_0 g^n$ we obtain from Eq. (2..1)

$$E(D_{n+1}|D_n) = \left(2 - \frac{D_n}{N_0} \right) D_n. \tag{2..3}$$

The growth factor g is eliminated and therefore N_0 is the only relevant quantity. Not only that, but the free of g property holds for the general case where g depends on n and then g_n is the growth factor from the generation n to $(n+1)$. To realize that, we replace in the former procedure:

$$g \rightarrow g_n, \quad g^n \rightarrow (g_0 \cdot g_1 \cdot g_2 \dots \cdot g_{n-1}) = \prod_{i=0}^{n-1} g_i. \tag{2..4}$$

Then $N_n = N_0 \cdot \prod_{i=0}^{n-1} g_i$. Usually $g_n \geq 1$, but at least we need for the first few generations $g_n > 1/2$ to avoid extinction of D_n (discussed in Section 6), and for later generations g_n to be bounded away from zero. With the normalization:

$$D_n = \tilde{D}_n / \prod_{i=0}^{n-1} g_i, \quad C_n = \tilde{C}_n / \prod_{i=0}^{n-1} g_i. \tag{2..5}$$

Eq. (2..1) with $g \rightarrow g_n$ transforms to Eq. (2..3) which is free of all g_n .

In section 3 we view this statistical process as a simple random walk model. We will show that $E(D_n^2) \approx E^2(D_n)$, and therefore the recursive Eq. (2..3) can be written as:

$$E(D_{n+1}) \approx \left(2 - \frac{ED_n}{N_0}\right) ED_n. \quad (2..6)$$

The Second Presentation: We split the population into males and females, $N_n^m = N_n^f = \frac{1}{2}N_n$, $D_n^m = D_n^f = \frac{1}{2}D_n$, $C_n^m = C_n^f = \frac{1}{2}C_n$ where “m” and “f” stand respectively for male and female for the D_n (Dvidians) and the rest $C_n = N_0 - D_n$.

We continue to use the normalized population numbers D_n , C_n and N_0 eliminating g from the equations since it doesn't affect the distribution.

Then the normalized population ratios are:

$$\begin{aligned} \frac{D_n^m}{N_0^m} &= \frac{D_n^f}{N_0^f} = \frac{D_n}{N_0} = \frac{\tilde{D}_n}{N_n} \equiv r_n \\ \frac{C_n^m}{N_0^m} &= \frac{C_n^f}{N_0^f} = \frac{C_n}{N_0} = \frac{\tilde{C}_n}{N_n} = \frac{N_0 - D_n}{N_0} = 1 - r_n. \end{aligned} \quad (2..7)$$

Now we sort the couple types and count them with their probabilities. (One can envision it again by the two card boxes described above or by a Roulette wheel procedure: The D_n^m and C_n^m are randomly marked in the cycling wheel and the D_n^f and C_n^f are randomly marked in the stationary platform.)

The couple types and their probabilities are:

$$\begin{aligned} (1) \quad D_n^m &\rightarrow D_n^f : D_n^m r_n = \frac{N_0}{2} r_n^2 \\ (2) \quad D_n^f &\rightarrow C_n^f : D_n^m (1 - r_n) = \frac{N_0}{2} r_n (1 - r_n) \\ (3) \quad C_n^m &\rightarrow D_n^f : C_n^m r_n = \frac{N_0}{2} (1 - r_n) r_n \\ (4) \quad C_n^m &\rightarrow C_n^f : C_n^m (1 - r_n) = \frac{N_0}{2} (1 - r_n)^2 \end{aligned} \quad (2..8)$$

The first three types give for each couple two D-children, and the fourth one gives zero

D-children, giving altogether for the next generation:

$$D_{n+1} = N_0(2 - r_n)r_n = \left(2 - \frac{D_n}{N_0}\right) D_n$$

which is the same equation as Eq. (2..3).

Upper bound estimate:

We would have liked to solve (2..3) recursively for ED_n up to $n = N$. This however cannot be done since:

$$E(D_n^2) = (ED_n)^2 + E(D_n - ED_n)^2. \quad (2..9)$$

and hence (Jensen's inequality)

$$ED_n^2 \geq (ED_n)^2.$$

The same holds for conditional expectations. Then by Eq. (2..3):

$$E(D_{n+1}) \leq 2ED_n - \frac{E^2(D_n)}{N_0}. \quad (2..10)$$

Therefore, we can solve recursively the equation:

$$E \hat{D}_{n+1} = 2 E \hat{D}_n - \frac{(E \hat{D}_n)^2}{N_0} \quad (2..11)$$

but the solution will be an *upper bound* to the solution of (2..3) for ED_n .

In the next section we will show that $(ED_n)^2$ is a very good approximation to $E(D_n^2)$ and consequently Eq. (2..6) will be a very good approximation for $N_0 \gg 1$.

3. The random walk model for $E[(D_{n+1} - D_n)|D_n]$

We wish to evaluate the standard deviation of our statistical process at the n th generation that leads to the descendant population of the $(n+1)$ generation. It is needed for evaluating

Eq. (2..9), and for that we apply a random walk model [2, 4].

Consider the unordered collection of the pairs of cards described in section 2, but without those cards where both couple members are non-descendants. It will yield D_n pairs. Next arrange these D_n pairs randomly and independently as a sequence running from 1 to D_n . Let i , $1 \leq i \leq D_n$ denote this sequence. Card i can have on it a pair of Davidians or one Davidian and one non-Davidian. Set $\lambda(i) = 2$ in case that a Davidian married a non-Davidian and $\lambda(i) = 1$ if Davidian married a Davidian. The sequence $\lambda(i), i = 1, 2, \dots, D_n$ constitutes a random walk process of D_n moves with a probability (D_n/N_n) to make 1 step ($\times 1$ D descendants), and a probability $(1 - D_n/N_n)$ to make 2 steps ($\times 2$ non-D descendants). The total average distance of the walk gives of next generation Davidian descendants D_{n+1} , given by Eq. (2..3). A similar procedure can be attributed for $D_{n+1} - D_n$ (successive years difference) obtained from Eq. (2..3):

$$\begin{aligned} E[(D_{n+1} - D_n)|D_n] &= \left(1 - \frac{D_n}{N_0}\right) D_n \\ &= \left[1 \cdot \left(1 - \frac{D_n}{2N_0}\right) + (-1) \cdot \frac{D_n}{2N_0}\right] D_n. \end{aligned} \quad (3..1)$$

Here the process is mapped to the standard random walk model where each move is of ± 1 (one forward or one backward) step. The total number of moves is D_n , each is either a unit step to the right with a probability $p = (1 - D_n/2N_0)$ or a unit step to the left with a probability $q = D_n/2N_0$.

The standard deviation of $(D_{n+1} - D_n)$ of this random walk process is known to be given by [2, 4]:

$$\begin{aligned} E^{1/2} \left\{ \left[(D_{n+1} - D_n) - E[(D_{n+1} - D_n)|D_n] \right]^2 | D_n \right\} &= \sqrt{D_n p q} \\ &= \sqrt{D_n \left(1 - \frac{D_n}{2N_0}\right) \left(\frac{D_n}{2N_0}\right)} ; \end{aligned} \quad (3..2)$$

but the left hand side of this equation is equal to:

$$E^{1/2} \left\{ \left[D_{n+1} - E(D_{n+1}|D_n) \right]^2 | D_n \right\} \quad (3.3)$$

which is the standard deviation of D_{n+1} .

Therefore:

$$\frac{E^{1/2} \left\{ \left[D_{n+1} - E(D_{n+1}|D_n) \right]^2 | D_n \right\}}{E(D_{n+1}|D_n)} = \frac{\sqrt{D_n \left(1 - \frac{D_n}{2N_0} \right) \left(\frac{D_n}{2N_0} \right)}}{\left(2 - \frac{D_n}{N_0} \right) D_n} = \sqrt{\frac{c}{N_0}}, \quad (3.4)$$

where $c \in (1/8, 1/4]$ is a number of order 1.

Then from Eq. (2.9) we have:

$$E(D_{n+1}^2|D_n) = E^2(D_{n+1}|D_n) \left(1 + \frac{c}{N_0} \right), \quad (3.5)$$

valid for any n . Therefore $E^2 D_n \approx E D_n^2$ and we will have for the recursive equation (2.3):

$$E(D_{n+1}) \approx \left(2 - \frac{E D_n}{N_0} \right) E D_n. \quad (3.6)$$

Then the descendant ratio $r_n = \frac{E \tilde{D}_n}{N_n} = \frac{E D_n}{N_0}$ (and $r_0 = \frac{1}{N_0}$) will be given by:

$$r_{n+1} \approx (2 - r_n) r_n \quad (3.7)$$

The last two equation are **key results** from which we derive our main conclusions.

It should be noted that the accumulated error in the recursive process along n (in our cases $n = 1, 2, \dots, 50$), is negligible since the relative error we found for a step in n is very small, of the order of $1/N_0^{1/2}$ (for any n), and $N_0 \sim 10^3 - 10^9$.

4. Examples for the recursive equation solution

We give below examples with plots of the descendant population ratio $r_n \approx \frac{E(\tilde{D}_n)}{N_n} = \frac{E(D_n)}{N_0}$ along the generation number, given by the exact solution of the recursive equation (3.7): $r_{n+1} \approx (2 - r_n)r_n$.

The plots in Fig. 1 give the r_n dependance on n for various values of N_0 (corresponding to various $r_0 = 1/N_0$). For a population of 10^5 , we get for the DAD full spreading time (the number of generations for the whole population to becomes descendants) about 20 (or $20 \times 25 = 500$ years). For each additional factor of 10 in N_0 we need $1/\log_{10} 2 = 3.22$ more generations. Thus even for a population of $N_0 = 10^9$ it takes only $(9/\log_{10} 2) + 4 = 33.9$ generations (850 years) to the DAD full spreading time.




Figure 1: Descendant population ratio r_n vs. n , obtained from the recursive equation - Eq. (3.7), for various starting population numbers: $N_0 = 1/r_0 = 10^3, 10^5, 10^7$ and 10^9 , (corresponding to plots from left to right, respectively).

5. Approximating the recursive equation with a differential equation

We derive from the recursive equation (3.6) an approximated differential equation. It will have the advantage that it possess an explicit solution, although the fitting is not exact.

We have from Eq. (3.6):

$$ED_{n+1} - ED_n \approx \left(1 - \frac{ED_n}{N_0}\right) ED_n. \quad (5.1)$$

Replacing ED_n with a continuous and differentiable function $\varphi(n)$, we obtain the equation:

$$\frac{d\varphi(n)}{dn} = \left(1 - \frac{\varphi(n)}{N_0}\right) \varphi(n). \quad (5.2)$$

Hence

$$dn = \frac{d\varphi(n)}{\varphi(n) \left(1 - \frac{\varphi(n)}{N_0}\right)}. \quad (5.3)$$

Set $r = \frac{\varphi(n)}{N_0}$, $r \in (0, 1)$ and $r_0 = 1/N_0$, then

$$\int_{r_0}^r \frac{dr}{r(1-r)} = \int_0^n dn. \quad (5.4)$$

Hence

$$n = \ln \frac{r/r_0}{(1-r)/(1-r_0)} = \ln \left(\frac{N_0 r \left(1 - \frac{1}{N_0}\right)}{1-r} \right). \quad (5.5)$$

Solving for r yields

$$r(n) = \frac{e^n}{e^n + N_0 - 1}. \quad (5.6)$$

This solution is only a rough estimate for the recursive equation - Eq. (3.7), and deviates from its exact solution especially for low n including the transition region to $r \rightarrow 1$. By

rescaling $n \rightarrow n \ln 2$, we obtain a modified equation:

$$r(n) = \frac{2^n}{2^n + N_0 - 1} \quad , \quad (5..7)$$

that describes the initial descendant evolution more accurately. However, it increases the width of the transition region. Eqs. (5..6) and (5..7) provide a rule of thumb approximation that $r(n) = 1/2$ is obtained at $2^n \approx N_0$.

Fig. 2 shows plots of the solutions - Eqs. (5..6) and (5..7) of the two differential equations which are compared to the exact solution given by the recursive equation - Eq. (3..7).




Figure 2: Descendant population ratio r_n obtained from: (a) the recursive equation (3..7), and the approximations (b) Eq. (5..6), and (c) Eq. (5..7)

6. Dynasty Extinction

Our model uses a growth factor that can vary from generation to generation, but takes a fixed average number of children of $m (= 2g)$ per family for the whole population. The overall DAD model that we presented gives an accurate description of the descendant distribution, but as we said, it needs attention to the first few generations. Then, when the dynasty starts to build up (D_n is a small number) it has fragility features just as the Galton-Watson (GW) process [1, 2]. The GW process was presented to find the stability and the extinction of surnames (or families) that is dependent on the average number of sons (or children) per family. We already mentioned that we need $m > 1$ ($g > 1/2$), just as in the GW process, to eliminate the possibility for the D extinction. However even for larger g there is a certain nonzero probability that a dynasty disappears. We need for that the probabilities for the various number of children per two parents that includes a finite probability for having no children. The zero children probability causes the extinction. We give here a brief analysis that basically follows the GW process.

For a Poissonic distribution the probability for the number of children k per family is $p_k = m^k e^{-m} / k!$. For example, for $m = 2$ ($g = 1$): $p_0 = 0.135$, $p_1 = 0.27$, $p_2 = 0.27$, and $p_3 = 0.18$. It is easy to realize that most of the extinction probability comes from a descendant chain of successive families along the generation line n , each with one child, terminated by a zero children family. The probability for that is the sum of the geometric series $\sum_n p_1^n p_0 = 0.185$. For $m = 3$ the sum is 0.058. We will see below that these numbers are very close to the results obtained by the GW calculation. In the GW process for Poissonic offspring distribution, the extinction probability can be obtained from the probability generating function [2]:

$$\tilde{s}_{n+1} = e^{m(\tilde{s}_n - 1)}, \quad (6.1)$$

and the survival probability $s_n \equiv 1 - \tilde{s}_n$ is given by:

$$s_{n+1} = 1 - e^{-ms_n}. \quad (6.2)$$

For $m \leq 1$ ($g \leq 1/2$) this recursion levels off to zero (i.e. family extinction). For $m > 1$ it levels off to a nonzero value given by the nontrivial fixed point of Eq. (6.2): one of the two roots of $s = 1 - e^{-ms}$. The trivial fixed point is always $s_a = 0$, and the second

one s_b , that depends on m , gives the survival probability. [When starting the recursion at $s_0 = 1/N_0$ it gives for this process the evolution along n of the descendant population ratio that asymptotically reaches s_b .] For $m = 2$ ($g = 1$) children per family, the survival ratio is $s_b = 0.797$ (extinction ratio of 0.203). For $m = 3$ ($g = 1.5$), the survival ratio is 0.94 (extinction ratio of 0.06) and for $m = 4$ ($g = 2$) it is $0.99 = 99\%$ (extinction ratio of $0.01 = 1\%$). The value of m can vary along the time and from place to place; the long term global population growth factor, however, is not much higher than $g = 1$. For $g \in (1, 1.2)$, i.e. $m \in (2, 2.4)$, the global family survival ratio is $s_b \in (0.8, 0.88)$.

The conclusion is that a certain fraction $(1 - s_b)$ of the population at the $n = 0$ generation will eventually have no descendants. The extinction that can occur in the first few generations is related to the GW process. We note what is discussed in the next section, that the extinction part doesn't affect the basic DAD process that is governed by the average g for those (the most) who survive. Nevertheless, it means about being a descendant of King David (or of anybody else) that: *either all Israelites are his descendant or there isn't even a single descendant*; or: *if there is for sure one descendant, then the whole population are descendants*. We said that for a moderate population growth the extinction part is $(10 - 20)\%$, meaning that there are no descendants for that fraction of the ancestors population. Nevertheless, as we said in the introduction, it is very likely that King David's dynasty survived the first few generations and then the DAD process ensured that they fully spread to the whole population.

7. Forward vs. backward approaches; Descendants vs. ancestors distributions

Our model was constructed by going from an ancestor to his descendants and thus look at the descendant distribution. Then it is possible to follow the path down from generation to generation, but the number of children per couple can vary and have some average value of $2g$. The Galton-Watson (GW) process [1, 2] is an example to such analysis. We note the basic difference between the GW process and ours. The DAD process depends on the question to

whom a Davidian is married. A non-Davidian mate contributes $m (= 2g)$ children per two parents, while an intra tribe marriage (Davidian to Davidian) gives only half of it, $m/2 = g$ children. In the GW like process this question is irrelevant. Its focus is on the distribution of the number of children per family that affect the descendant statistics, but it doesn't include the interplay between intra and inter tribe marriage. Therefore the GW process is relevant in the DAD analysis in the first generations only to understand the fragility and extinction. At that stage the DAD process is anyhow similar to the GW one since the probability for a non-Davidian mate is almost 100%. This is the reason why we could use in Section 6 the GW analysis for evaluating the survival ratio.

It is also possible to go along the opposite direction from a descendant back in time to his ancestors. Then the route follows the ancestors distribution rather than the descendants. The backward way is more common because of the simplicity to track genealogical family trees of more recent generations, and since then in the analysis the number of parents is a fixed value of 2. Nevertheless the different backward trajectories that can pass through the same ancestors have probabilistic aspects. This viewpoint alludes on another aspect, the number of paths that connect a descendant at the n -th generation to a specific ancestor. This issue depends on the growth factor and the degrees of inter versus intra community mobility. It is more likely that the connection path number is significantly higher within communities with a common social-geographic history. Derrida et al. [5, 6] took the backward direction analysis and obtained much understanding on various sides of the process. We mention here only one of their results on the population ratio of ancestors at a backward generations given in Refs. [5, 6]. The result there is related to the GW process giving for the non-ancestors population ratio at a backward generation i the recursive equation: $\tilde{s}_{i+1} = e^{m(\tilde{s}_i-1)}$. For the ancestor ratio $s_i \equiv 1 - \tilde{s}_i$, it becomes $s_{i+1} = 1 - e^{-ms_i}$ (dependent on m or g), which is equivalent to Eq. (6.2) of the GW process, and so it differs from our approach.

8. Conclusions

We have presented a model for the distribution of descendants along generations, the DAD route. The descendant population and the ratio are given by the recursive equations (3.6) and (3.7). The descendant ratio, given for a few examples in figure 1, is shown to reach 1 in a relatively few generations. For an initial population of N_0 the DAD full spreading time is $\sim \log N_0 / \log 2 + 4$, that gives about 20 generations (500 years) for $N_0 = 10^5$. Every additional factor of 10 in N_0 adds $1/\log_{10} 2 = 3.32$ generations to the DAD full spreading time. The basic DAD route behavior, in particular the descendant population ratio does not depend on the population growth factor g , but only on the initial population N_0 .

We have not included here any genetic or geo-social-demographic aspects. It is clear that DAD will not spread into and out of very isolated groups. Nevertheless, we saw how quick the spreading process is. For a small group of 10^3 or 10^4 it takes about 14 and 17 generations (350 and 425 years) to reach a full descendant spreading ratio. For all Israelites at King's David time it took 24-26 generations (600-650 years) to make every Israelite his descendent. That therefore happened already at 400 BC, the beginning of the era of the Second Temple in Jerusalem. Even for the whole world population at King's David time (1000 BC), estimated as 5×10^7 , it is but 29.5 generations (740 years). Segregation of local communities can slow down the process, but only in a limited way for relatively short time. It would be sufficient that one descendant migrates to another community, say with a similar population number, to make the whole population descendants in a few generations. Therefore more globally one might say that even if mankind started with many Adams and Eves, it took relatively very short time to consider the whole founder group at any early era, as a Common Forefather – Super DAD – of all of us.

For the future, the DAD route means that, assuming a reasonable population mobility, each of us on earth today ($N_0 \approx 5 \times 10^9$) (beyond the low extinction percentage discussed in Section 6, and assuming that no catastrophic event happens, will be an ancestor of everyone in the world - on Earth and beyond?) in $\sim \log(5 \times 10^9) / \log 2 + 4 \approx 36$ generations (~ 900 years) from now. Can this picture lend a philosophical meaning to what is said about a

common forefather of mankind? We saw that all of us have common ancestors and eventually we will be ourselves the ancestors of everyone in the future in a relatively short period of time. They all were our Fathers and Mothers and they all will be our Sons Daughters...

We finally note that after we first deposited the paper in ArXiv, our attention was drawn to the work of Derrida et al. on the same subject [5, 6]. We believe that our paper adds new insight and results treating the process along the generation line rather than the backward way. We have added a short discussion on the difference between the approaches in Sec. 7.

Acknowledgement: We wish to thank Israel Bar-David, Nahum Shimkin, Adam Shwartz, and Ofer Zeitouni for useful remarks. We also thank Yosi Avron and Yosef Maruvka for drawing our attention to the work of Bernard Derrida et al. on the subject.

References

- [1] F. Galton and H.W. Watson, “On the probability of extinction of families”, Journal of the Anthropological Institute, (1874), 308–311.
- [2] S. Karlin and H. H. Taylor, “A First Course in Stochastic Processes”, Academic Press, New York, 1975.
- [3] The Bible, I Chronicles; Chapter 21:5, (New American Standard Bible, 1995): “Joab gave the number of the census of all the people to David. And all Israel were 1,100,000 men who drew the sword; and Judah was 470,000 men who drew the sword.” We therefore estimate the total population to be about three times larger - 4 to 5 million.
- [4] Wolfram Math World on “Random walk - 1-Dimensional”,
<http://mathworld.wolfram.com/RandomWalk1-Dimensional.html>
- [5] B. Derrida, S. C. Manrubia, and D. H. Zanette; “Statistical Properties of Genealogical Trees”, Phys. Rev. Lett. 82, 1987, (1999).
- [6] B. Derrida, S.C. Manrubia, D.H. Zanette, Jour. theor. Biol. 203 (2000) 303.
(arXiv:physics/0003016v1); <http://arxiv.org/abs/physics/0003016>